# The Effects of Exploitation on the Arctic Charr Population of the Sylvia Grinnell River, Northwest Territories

A. H. Kristofferson and R. D. Sopuck

Western Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

November 1983

**Canadian Manuscript Report of Fisheries & Aquatic Sciences No. 1721**

2008-06002/

## Canadian Manuscript Report of Fisheries and Aquatic Sciences

These reports contain scientific and technical information that represents an important contribution to existing knowledge but which for some reason may not be appropriate for primary scientific (i.e. *journal*) publication. They differ from Technical Reports in terms of subject scope and potential audience: Manuscript Reports deal primarily with national or regional problems and distribution is generally restricted to institutions or individuals located in particular regions of Canada. No restriction is placed on subject matter and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries management, technology and development, ocean sciences, and aquatic environments relevant to Canada.

Manuscript Reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report will be abstracted by *Aquatic Sciences and Fisheries Abstracts* and will be indexed annually in the Department's index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 901-1425 were issued as Manuscript Reports of the Fisheries Research Board of Canada. Numbers 1426-1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Details on the availability of Manuscript Reports in hard copy may be obtained from the issuing establishment indicated on the front cover.

## Rapport manuscrit canadien des sciences halieutiques et aquatiques

Ces rapports contiennent des renseignements scientifiques et techniques qui constituent une contribution importante aux connaissances actuelles mais qui, pour une raison ou pour une autre, ne semblent pas appropriés pour la publication dans un journal scientifique. Ils se distinguent des Rapports techniques par la portée du sujet et le lecteur visé; en effet, ils s'attachent principalement à des problèmes d'ordre national ou régional et la distribution en est généralement limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du Ministère des Pêches et des Océans, notamment gestion des pêches; techniques et développement, sciences océaniques et environnements aquatiques, au Canada.

Les Manuscrits peuvent être considérés comme des publications complètes. Le titre exact paraît au haut du résumé de chaque rapport, qui sera publié dans la revue *Aquatic Sciences and Fisheries Abstracts* et qui figuera dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros de 1 à 900 de cette série ont été publiés à titre de manuscrits (Série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés en tant que manuscrits (Série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros allant de 901 à 1425 ont été publiés à titre de manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 ont été publiés à titre de Rapport manuscrits du Service des pêches et de la mer, Ministère des Pêches et de l'Environnement. Le nom de la série a été changé à partir du rapport numéro 1551.

La page couverture porte le nom de l'établissement auteur où l'on peut se procurer les rapports sous couverture cartonnée.

Canadian Manuscript Report of
Fisheries and Aquatic Sciences 1721

November 1983

# THE EFFECTS OF EXPLOITATION ON THE ARCTIC CHARR POPULATION OF THE SYLVIA GRINNELL RIVER, NORTHWEST TERRITORIES

by

A. H. Kristofferson and R. D. Sopuck

Western Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

This is the 44th Manuscript Report
from the Western Region, Winnipeg

© Minister of Supply and Services Canada 1983

Cat. no. 97-4/1721 ISSN 0706-6473

Correct citation for this publication is:

KRISTOFFERSON, A.H., and R.D. SOPUCK. 1983. The effects of exploitation on the Arctic charr population of the Sylvia Grinnell River, Northwest Territories. Can. Manuscr. Rep. Fish. Aquat. Sci. 1721: v + 35 p.

TABLE OF CONTENTS


| | Page |
|---|---|
| ABSTRACT/RESUME | v |
| INTRODUCTION | 1 |
| MATERIALS AND METHODS | 1 |
| Study area | 1 |
| Creel census (sport fishery) | 1 |
| Experimental gillnetting | 1 |
| Subsistence fishery | 1 |
| Biological data | 1 |
| Data analysis | 2 |
| RESULTS AND DISCUSSION | 2 |
| Status of the stock | 2 |
| Length- and age-frequency distribution | 2 |
| Mortality | 2 |
| Growth | 3 |
| Maturity and spawning potential | 3 |
| Sport fishery | 4 |
| Subsistence fishery | 4 |
| Sylvia Grinnell Lake | 5 |
| Effect of fisheries on the charr stock | 5 |
| Recovery of the charr stock | 5 |
| RECOMMENDATIONS | 6 |
| ACKNOWLEDGMENTS | 6 |
| REFERENCES | 7 |


LIST OF FIGURES


| Figure | | Page |
|---|---|---|
| 1 | Map of the Frobisher Bay area, Baffin Island, showing the Sylvia Grinnell River | 8 |
| 2 | Map of the outlet of the Sylvia Grinnell River showing the study area | 9 |
| 3 | A comparison of length-frequency distributions of samples of Arctic charr taken from the Sylvia Grinnell River during 1948/50/51, 1958 and 1976/77 | 10 |
| 4 | A comparison of age-frequency distributions of samples of Arctic charr taken from the Sylvia Grinnell River during 1948/50/51 and 1976/77 | 10 |
| 5 | Catch curves for Arctic charr taken from the Sylvia Grinnell River during 1948/50/51, 1958 and 1976/77 | 11 |
| 6 | A comparison of mortality rates (catch curve) for Arctic charr, aged 13 (14) to 18 years, taken from the Sylvia Grinnell River during 1948/50/51, 1958 and 1976/77 | 11 |
| 7 | A comparison of growth rates of Arctic charr taken from the Sylvia Grinnell River in 1948/50/51 and 1976/77 with those of Arctic charr from other rivers in the Northwest Territories | 12 |
| 8 | A comparison of length- and age-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by angling in 1976 and 1977 | 13 |
| 9 | A comparison of length- and age-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by experimental gillnet in 1976 and 1977 | 14 |
| 10 | Length-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by experimental gillnets on four occasions during 1977 | 15 |
| 11 | Daily catch of Arctic charr from the Sylvia Grinnell River taken by angling and snagging compared with daily high tide level, 1976 | 16 |
| 12 | Daily catch of Arctic charr from the Sylvia Grinnell River taken by angling and snagging compared with daily high tide level, 1977 | 16 |
| 13 | A comparison of length-frequency distributions of Arctic charr taken by angling from the Sylvia Grinnell River in 1976/77, the Tree River in 1978 and the Robertson River in 1979 | 17 |
| 14 | A comparison of length-frequency distributions of Arctic charr taken by different types of gear from the Sylvia Grinnell River, 1976/77 | 17 |
| 15 | A comparison of length-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by snagging in 1976 and 1977 | 18 |
| 16 | Length-frequency distribution of Arctic charr taken from Sylvia Grinnell Lake by experimental gillnet in 1977 | 18 |
| 17 | Length- and age-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by experimental gillnet in 1982 | 19 |


LIST OF TABLES


| Table | | Page |
|---|---|---|
| 1 | Instantaneous total and fishing mortality and annual survival of | |

| Table | | Page |
|---|---|---|
| | Arctic charr, aged 13 to 18 years, from the Sylvia Grinnell River in 1948/50/51, 1958 and 1976/77 | 20 |
| 2 | Weight-length relationships of Arctic charr from a number of locations compared with those from the Sylvia Grinnell River, 1976 and 1977 | 20 |
| 3 | Condition factor (K), of Arctic charr caught in the Sylvia Grinnell River in 1976 and 1977 | 21 |
| 4 | Age, mean length and size range of Arctic charr from the Sylvia Grinnell River, 1948, 1950 and 1951 | 22 |
| 5 | Percent of Arctic charr, by length interval, captured in spawning condition while migrating up the Sylvia Grinnell River in 1976/77 | 23 |
| 6 | Percent of Arctic charr, by age, captured in spawning condition while migrating up the Sylvia Grinnell River in 1976/77 | 24 |
| 7 | A comparison of creel census data for Arctic charr angled in the Sylvia Grinnell River in 1958, 1976/77, the Tree River in 1978 and the Robertson River in 1979 | 25 |
| 8 | Catch record of the subsistence gillnet fishery in the Sylvia Grinnell River, August 1977 | 26 |
| 9 | Annual harvest statistics for Arctic charr from the Sylvia Grinnell River, 1958-66 | 27 |
| 10 | Annual harvest, mean round weight per charr and number of charr taken by fisheries at the Sylvia Grinnell River in 1948, 1950, 1958 and 1977 | 28 |
| 11 | Number of days fished, harvest, mean round weight per charr and number of charr taken by angling, subsistence gillnetting and snagging at the Sylvia Grinnell River, 1977 | 28 |


LIST OF APPENDICES


| Appendix | | Page |
|---|---|---|
| 1 | Summary of biological data by length interval for Arctic charr taken by angling in the Sylvia Grinnell River, 1976 | 29 |
| 2 | Summary of biological data by age for Arctic charr taken by angling in the Sylvia Grinnell River, 1976 | 29 |
| 3 | Summary of biological data by length interval for Arctic charr taken by angling in the Sylvia Grinnell River, 1977 | 30 |
| 4 | Summary of biological data by age for Arctic charr taken by angling in the Sylvia Grinnell River, 1977 | 30 |
| 5 | Summary of biological data by length interval for Arctic charr taken by experimental gillnets in the Sylvia Grinnell River, 1976 | 31 |
| 6 | Summary of biological data by age for Arctic charr taken by experimental gillnets in the Sylvia Grinnell River, 1976 | 31 |
| 7 | Summary of biological data by length interval for Arctic charr taken by experimental gillnets in the Sylvia Grinnell River, 1977 | 32 |
| 8 | Summary of biological data by age for Arctic charr taken by experimental gillnets in the Sylvia Grinnell River, 1977 | 32 |
| 9 | Summary of biological data by length interval for Arctic charr taken by subsistence gillnets in the Sylvia Grinnell River, 1977 | 33 |
| 10 | Summary of biological data by age for Arctic charr taken by subsistence gillnets in the Sylvia Grinnell River, 1977 | 33 |
| 11 | Summary of biological data by length interval for Arctic charr taken by snagging in the Sylvia Grinnell River, 1977 | 34 |
| 12 | Summary of biological data by age for Arctic charr taken by snagging in the Sylvia Grinnell River, 1977 | 34 |
| 13 | Description of the relative stages of maturity of Arctic charr based on gross examination of the gonads | 35 |

## ABSTRACT

KRISTOFFERSON, A.H., and R.D. SOPUCK. 1983. The effects of exploitation on the Arctic charr population of the Sylvia Grinnell River, Northwest Territories. Can. Manuscr. Rep. Fish. Aquat. Sci. 1721: v + 35 p.

A creel census and biological investigation of the Arctic charr (Salvelinus alpinus) population of the Sylvia Grinnell River was conducted in July and August of 1976 and 1977. Four hundred and twenty-seven anglers interviewed in 1976 retained a total of 212 charr and 374 interviewed in 1977 retained a total of 379 charr. Estimates suggest that the subsistence gillnet fishery took at least 2 282 charr in 1977 and snagging took at least 4 923 charr in 1977. The size and age composition of the stock has declined considerably since earlier studies in 1948, 1950, 1951 and 1958. Charr larger than 55 cm were rarely encountered during this study. Observations of the 1982 fishery indicate that the stock has not yet recovered. Present fishing effort is probably sufficient to prevent the recovery of the stock. Recommendations include closure of a portion of the river below the falls to all fishing since charr congregate there in large numbers and are very vulnerable to capture. Such a closure would eliminate snagging, a practice which harvests large numbers of small charr.



Key words: age composition; angling; catching methods; fishing mortality; monitoring; population structure; Salvelinus alpinus (L); size distribution.



## RESUME

KRISTOFFERSON, A.H., and R.D. SOPUCK. 1983. The effects of exploitation on the Arctic charr population of the Sylvia Grinnell River, Northwest Territories. Can. Manuscr. Rep. Fish. Aquat. Sci. 1721: v + 35 p.

En juillet et août 1976 et 1977, on a effectué un relevé des prises d'ombles chevaliers (Salvelinus alpinus) ainsi qu'une étude biologique des stocks d'ombles de la rivière Sylvia Grinnell. Les 427 pêcheurs sportifs interviewés en 1976 ont gardé un total de 212 ombles, tandis que les 374 pêcheurs interviewés en 1977 ont gardé 379 ombles. On estime que la pêche de subsistance aux filets maillants a récolté 2 282 ombles en 1977 et qu'au moins 4 923 ombles ont été pris à l'hameçon nu en 1977. La taille et l'âge moyens des poissons ont baissé considérablement depuis les études de 1948, 1950, 1951 et 1958; lors de la présente étude, les ombles de plus de 55 cm étaient rares. Une étude des pêches de 1982 a révélé que les stocks d'ombles n'étaient pas encore rétablis. La pêche actuelle est probablement suffisante pour empêcher ce rétablissement. Cet ouvrage recommande d'interdire la pêche dans la partie de la rivière, en aval des chutes, où les poissons se rassemblent en grand nombre et sont très faciles à prendre. La prise d'une telle mesure éliminerait la pêche à l'hameçon nu, méthode qui récolte un grandnombre de petits ombles.



Mots-clés: moyenne d'âge; pêche sportive; méthodes de pêche; mortalité due à la pêche; contrôle; structure de la population; Salvelinus alpinus (L); distribution des tailles.

## INTRODUCTION

The Arctic charr (Salvelinus alpinus (L.)) of the Sylvia Grinnell River on Baffin Island have been subject to varying degrees of exploitation by man. Inuit have traditionally fished for charr in the lower reaches of this river where it empties into Frobisher Bay (Hunter 1976). Commercial fishing using gillnets was initiated by the Shaw Steamship Company in 1947 and continued in 1948 and 1950 (Hunter 1976). Fishing was done in the estuary adjacent to the mouth of the river (Grainger 1953). Some of the older residents of Frobisher Bay claim that this fishery removed large quantities of charr each year. During 1951-57 charr were taken by subsistence and recreational fisheries (Hunter 1976). In 1958 the Department of Indian Affairs and Northern Development reestablished the commercial fishery for charr with an annual harvest objective of 4 500 kg (Hunter 1976). During 1958-62 fishing was good, with the quota being taken each year in only a few days (Hunter 1963). However, declining catch for effort and small fish size followed and the commercial fishery ceased operating after 1966 (Hunter 1976). Since then only the subsistence and recreational fisheries have persisted.

There are indications that Arctic charr in the Sylvia Grinnell River have been overexploited (Scott and Crossman 1973; Hunter 1976). Local residents and anglers in Frobisher Bay report that their catch consists mostly of small fish and return for fishing effort is poor. The present study was undertaken during the summers of 1976 and 1977 to assess the status of the Arctic charr stock in the river. Results are compared with those of previous studies by Grainger (1953) and Hunter (1958). Changes in the charr population over the period of study are discussed and recommendations are made for future use of this valuable resource.

## MATERIALS AND METHODS

### STUDY AREA

The Sylvia Grinnell River empties into Frobisher Bay at 63°44'N, 68°34'W (Fig. 1), about 2 km west of the town of Frobisher Bay. The river drains Sylvia Grinnell Lake, located approximately 70 km upstream. Anadromous Arctic charr are believed to overwinter in this lake (Grainger 1953). The river flows over a series of falls and rapids just before it meets tidal waters in an upper area of Frobisher Bay (Fig. 2). The height of the falls varies from zero to about 3.7 m depending on the tide. These falls present an impassable barrier to charr ascending the river except during spring tides when the falls are inundated by the high water (Grainger 1953). Spring tides occur about twice a month. During low water the charr congregate below the falls and it is here that most of the fishing takes place.

### CREEL CENSUS (SPORT FISHERY)

A creel census was conducted during the periods July 10 to August 28, 1976 and July 23 to August 15, 1977. Fishing sites (Fig. 2) were visited twice daily. As many anglers as possible were interviewed and their time spent fishing as well as the number of charr each one caught, kept and released was recorded. As many charr as possible from each angler's catch were sampled for biological data.

### EXPERIMENTAL GILLNETTING

Experimental gillnets were used to capture Arctic charr at the mouth of the river. The gillnets were made up of five panels, each 22.9 m long and each of a different mesh size as follows: 38, 63, 89, 114 and 139 mm, stretched measure. Overnight sets (18 h) were made on August 20 and 21, 1976. The nets were set in the channel with one end fastened to the shoreline. An overnight set (24 h) was made on July 28, 1977 followed by daytime sets (6 h each) on August 2, 12 and 15. In addition, two overnight floating sets (20 h each) were made in Sylvia Grinnell Lake on August 7 and 8, 1977. Charr captured by this method were sampled for biological data.

### SUBSISTENCE FISHERY

A subsistence gillnet fishery operated in the estuary during the 1976 and 1977 study periods. Some charr captured in this fishery on August 23 and 25, 1977 were sampled for biological data. Estimates of the daily catch were attempted in 1977 by observing the fishermen while they lifted their nets and removed the charr.

Another method of fishing was by snagging. Bare, weighted hooks were pulled through the water to hook fish. This was practised mostly by Inuit fishermen apparently for subsistence purposes. It was difficult to determine the number of charr taken by snagging since large numbers were captured in a short period of time by many individuals. Hence the number of charr recorded as having been taken by snagging is a minimum estimate.

### BIOLOGICAL DATA

Charr were sampled for fork length (±1 mm) and round weight (±25 g). Saggital otoliths were removed and stored dry in envelopes for age determination. The otoliths were then immersed in a 3:1 solution of benzyl-benzoate and methyl salicylate in the laboratory prior to aging. Some otoliths were ground slightly on a fine carborundum stone to improve annulus clarity. Annuli were counted while the otoliths were observed with the aid of a compound microscope (X20). Ages were determined following Grainger (1953), counting the central core of the otolith as the first annulus. Sex and maturity were determined by gross examination of the gonads. Maturity stages are described in Appendix 13.

Age- and length-frequency histograms were constructed to display catch compositions from various gears. Ages of charr younger than 11 years that were taken in 1958 were not available. Hence an age-frequency distribution was

not constructed, although a catch curve for those 11 years and older was. Weight-length relationships were calculated using least squares regression analysis on logarithmic transformations of fork lengths and round weights.

The relationship is described as:

$$\text{Log}_{10} W = a + b\ (\text{Log}_{10} L)$$

Where: W = weight in grams
L = fork length in millimetres

Mean fork length at age was plotted from the data and growth rate was determined visually. Relative condition factor (K), a measure of the plumpness or robustness of the fish, was determined as:

$$K = \frac{W \times 10^5}{L^3}$$

Where: W = weight in grams
L = fork length in millimetres

Instantaneous total mortality (Z) was calculated from least squares regression lines fitted to the descending limb of catch curves. Only that portion of the curve that appeared linear was included in the analysis. Annual survival rates were calculated from Z. Instantaneous natural mortality (M) was assumed to be 0.13 after Hunter (1958). Instantaneous fishing mortality (F) was calculated from Z = F + M.

## DATA ANALYSIS

Data were analyzed using a Hewlett Packard model 9810-A programmable calculator and an Amdahl 470/V7 Computer. Programs from the Statistical Analysis System (1979) were used for tests of significance and regression.

## RESULTS AND DISCUSSION

### STATUS OF THE STOCK

#### Length- and age-frequency distribution

Results of this study indicate that the composition of the anadromous Arctic charr stock in the Sylvia Grinnell River has changed significantly since 1948. Studies by Grainger (1953) show that charr larger than 55 cm (fork length) were relatively abundant in this system in 1948, 1950 and 1951. Data from Hunter (1958) reveal little change up to 1958. The present study indicates that charr larger than 55 cm had virtually disappeared from the stock by 1976/77 (Fig. 3). Correspondingly, a shift in the age composition of the charr stock is evident (Fig. 3). The gear used in each study are comparable. Grainger (1953) reports that gillnets of 25, 63, 102, 127, 139, 152 and 165 mm (stretched measure) were the primary method of capture. To a lesser extent hand nets, gaffs, jigs and minnow traps were used. Hunter (1958) used gillnets of 38, 51, 63, 89, 114 and 139 mm mesh size in 1958. During the present study samples were taken with similar sized gillnets (see MATERIALS AND METHODS) and by angling.

Johnson (1980) observed a bimodal length-frequency for anadromous Arctic charr taken in Creswell Bay in 1962. Subsequent examination of this population by Sekerak et al. (1976) and de March et al. (1978) revealed that charr in the larger mode were less frequent than in 1962. They attributed the decline in large charr to fishing pressure. Although bimodality in length of Arctic charr is not as pronounced in the Sylvia Grinnell River stock, there is some evidence of it in the 1948/50/51 and 1958 samples (Fig. 3). Johnson (1980) noted modal values of 35 cm and 55 cm in the Creswell Bay stock in 1962. A mode of 30 cm to 35 cm is evident in all three samples from the Sylvia Grinnell River (Fig. 3). Note that it is the only mode evident in the 1976/77 sample, the larger individuals in the population having been removed. Johnson (1980) maintains that the removal of the large population mode of Creswell Bay charr left only a small-sized, slow-growing group, with little capacity to yield a significant harvest. This appears to have happened to the Sylvia Grinnell River charr stock as well.

The modal age of charr in 1976/77 had fallen to 10 years from 13 years in 1948/50/51 (Fig. 4). By comparison to the marked reduction in modal length over this time period, the difference in model age seems relatively small. However, this would not be unexpected if this small mode was comprised of slow-growing individuals, many of which were relatively old for their size.

#### Mortality

The catch curve for 1948/50/51 has a gently sloping right-hand limb suggesting a relatively low mortality rate (Fig. 5). This is not surprising since commercial fishing was just beginning when the sample was taken. Examination of the curve from the 1958 data suggests mortality had increased only amongst the oldest charr. This could be attributable to the early commercial fishery. Age data for charr taken in 1958 were unavailable for those less than 11 years old, hence the left-hand limb of this curve is missing. The catch curve plotted from the 1976/77 data is markedly displaced to the left, indicating that the old charr have been removed from the stock.

A comparison of mortality rates of charr aged 13 to 18 years (14 to 18 years for 1958) is shown in Fig. 6. Instantaneous total mortality rates (Z) were calculated from the slopes of the straight lines fitted to the data. In each case the data show a good linear fit. These data indicate that instantaneous total mortality was relatively low in 1948/50/51 (Z = 0.31) and in 1958 (Z = 0.14). It had increased significantly (Z = 0.70) by 1976/77. If it is assumed that natural mortality has remained constant, the increase was due to increased fishing mortality (Table 1). Subsequently, annual survival (S) amongst these age groups dropped from 87% in 1958 to 50% in 1976/77 (Table 1). If natural mortality decreased, as it often does following exploitation (Tyler and Gallucci 1980), fishing mortality would be even higher than is calculated here.

## Growth

Weight-length relationships for Arctic charr taken in 1976 and 1977 (sexes combined) are as follows:

1976 $\quad Log_{10} W = -5.2101 + 3.0724\ Log_{10} L$
$\quad N = 255;\ SD_b = 0.0317$

1977 $\quad Log_{10} W = -5.3254 + 3.1368\ Log_{10} L$
$\quad N = 848;\ SD_b = 0.0259$

Using analysis of covariance, no significant differences ($p>0.05$) were found between sexes or between years (sexes combined) with respect to slopes or adjusted means.

Weight-length relationships of several charr stocks are shown in Table 2. Charr taken in the Sylvia Grinnell River show an increase in weight with length which is similar to other stocks. This increase appears to be isometric.

Mean condition factor, K, was lower in 1976 (K = 0.94, N = 255) than in 1977 (K = 1.06, N = 847) suggesting that 1976 was a poor growing year. Condition was lower over the entire range of sizes of charr sampled in 1976 compared with 1977 (Table 3). Johnson (1980) has observed that year to year variation in weight increase during the summer is quite high. The availability of food and time spent at sea no doubt contribute to this variation.

Charr from the Sylvia Grinnell River grow more slowly than those from the Robertson River on North Baffin Island, the Jayco River on Victoria Island, and the Tree River, on the mainland in the central Arctic (Fig. 7). All these charr stocks are subject to exploitation, with sport fisheries at the Robertson (Moshenko 1981) and Tree (Falk and Dahlke 1979) rivers and a commercial fishery at the Jayco River (Dahlke and Falk 1979). Tree River charr are well known for their rapid growth (Moshenko et al. 1983), as is the Sylvia Grinnell River stock for its slow growth (Grainger 1953). Also, there appears to have been an apparent decrease in growth rate of the Sylvia Grinnell stock by 1976/77 compared with the growth rate calculated by Grainger (1953) for 1948/50/51 (Fig. 7). Grainger's (1953) method of aging was used in the present study so this decrease is probably not due to a difference in aging technique.

As discussed earlier, Johnson (1980) found differences in growth rates between the large and small modes of charr investigated at Creswell Bay in 1962. Charr in the small mode ranged in age from 6 to 25 years while those in the large mode ranged from 11 to 28 years of age. Grainger (1953) expressed the growth rate of Sylvia Grinnell River charr as mean length at age. Differences in growth rates tend to be obscured when this technique is used (Johnson 1980). If the range in length at age from Grainger's (1953) sample is examined (Table 4) it can be seen that it is comprised of small, relatively old charr and large, relatively young charr. Charr aged 10 years ranged in length from 26.5 cm to 52 cm while those aged 17 years ranged from 36.5 cm to 70 cm in length. Therefore, the two size modes evident in Fig. 3 may well have been made up of charr with different growth rates. The growth rate presented by Grainger (1953) would be an average of the different growth rates, since mean length at age was used. Removal of the large, fast-growing mode by fishing would leave only the small, slow-growing mode resulting in the apparent decrease in growth rate observed by the present study (Fig. 7).

Johnson (1980) has observed bimodal size distributions in a number of Canadian Arctic charr stocks. However, the factors maintaining the segregation of the size groups are not known. Johnson (1976) has suggested that recruitment is controlled by density dependent factors in landlocked lakes. Amongst the anadromous stocks at Creswell Bay and Nauyuk Lake the density of the large stock does not appear to maintain the segregation between the two size groups. Johnson (1980) questions whether all members of the smaller mode, irrespective of age, are able to grow into the larger mode under appropriate conditions, or whether it becomes impossible after a certain age. Perhaps the size modes represent two separate forms of charr as advocated by a number of researchers (Johnson 1980).

The reduction in the density of the large modes at Creswell Bay, Nauyuk Lake and Sylvia Grinnell River appears to be related to exploitation. Continued, intense fishing pressure could be sufficient to prevent the return of the large size mode in these cases. Whether or not individuals in the small mode are capable of replenishing the larger size mode through accellerated growth will only be determined if fishing pressure is reduced.

## Maturity and spawning potential

Grainger (1953) concluded that female charr from the Sylvia Grinnell River first reached sexual maturity at a length of about 450 mm and an age of about 12 years. Once they reach maturity charr in most areas of the Northwest Territories do not spawn every year (Johnson 1980) and most remain in freshwater the summer prior to spawning (Sprules 1952; Grainger 1953; Johnson 1980). However, a few spawners do return from the sea (Moore 1975) as was observed during the present. Eight percent of females and one percent of males captured were in spawning condition. No males under 500 mm in fork length and no females under 350 mm in fork length were mature (Table 5). No males under 11 years of age and no females under 10 years of age were mature (Table 6). The length and age distributions of charr in the Sylvia Grinnell River in 1976/77 suggest then that there were few fish large enough and old enough to be considered potential spawners (Fig. 3 and 4). Although the relationship between recruitment and spawning stock size amongst anadromous Arctic charr is not well understood (Johnson 1980) it could be that a paucity of spawners in the Sylvia Grinnell River stock results in poor recruitment and thus inhibits productivity. Apparently the charr stock was in a reduced state in 1966 since the commercial fishery terminated at that time owing to declining catch per unit of effort. Hence the reduced status of the stock in 1976/77 could have been related to

poor recruitment from a reduced spawning stock.

Stocks under stress from a fishery are less able to survive environmental changes and even temporary changes could reduce or prevent their full recovery (Ricker 1977). The Sylvia Grinnell River charr stock, in its present reduced state, may now also be particularly vulnerable to environmental degration.

## SPORT FISHERY

An active sport fishery takes place at the Sylvia Grinnell River during summer. Four hundred and twenty-seven anglers interviewed spent a total of 871 h fishing in 1976 and 374 interviewed fished for 1 102 h in 1977 (Table 7). Obviously angling is an important recreational pastime in Frobisher Bay since 83% of those interviewed in 1977 were residents of the community. Mean length (36.4 cm, N = 137) and mean age (11.6 years, N = 123) of charr angled in 1976 was slightly larger and older respectively than in 1977 (33.8 cm, N = 259; 10.5 years, N = 218) (Fig. 8). It appears that the mean length (37.1 cm, N = 244) and age (11.3 years, N = 224) of charr available to anglers in 1977, as sampled by experimental gillnets (Fig. 9), was larger and older respectively than what the anglers took (Fig. 8). However, the converse was observed in 1976, where the mean fork length of charr taken by the experimental gillnets (28.8 cm, N = 118) (Fig. 9) was smaller and the mean age (9.2 years, N = 104) was younger than that taken by angling during that year (Fig. 8). However, the results of the experimental gillnetting in 1976 may not be representative, since it was carried out on only two consecutive days in late August. Angling took place throughout the duration of the study in 1976 and 1977 and experimental gillnetting in 1977 took place on four days over an 18 day period (Fig. 10). This figure indicates a gradual increase in the size of the charr in the river over the course of the summer, perhaps reflecting the return of migrants from the sea.

Most of the anglers interviewed were not successful at the Sylvia Grinnell River, but a small number of skillful anglers caught charr regularly. Their success appeared to be relatively constant throughout the 1976 season (Fig. 11). Angling success in 1977 showed a peak at the end of July (Fig. 12) coincident with the occurrence of the spring tide. Grainger (1953) observed that charr moved upstream to the falls on the incoming tides and then withdrew into the bay on the falling tide. This pattern is apparently repeated throughout the summer until the upstream overwintering migration takes place. Angling success is probably influenced by these movements.

In comparison to the quality of angling available at two well-known Arctic charr sport fisheries, namely the Tree and Robertson rivers, angling at the Sylvia Grinnell River is poor (Table 7). In 1978, 84% of anglers censused at the Tree River were successful in capturing at least one charr (Falk and Dahlke 1979) as were 95% at the Robertson River (Moshenko 1981) compared with 26% and 47% in 1976 and 1977, respectively, at the Sylvia Grinnell River. Mean fork length of charr taken at the Tree River was 75.6 cm and at the Robertson River 65.9 cm compared with 36.4 cm (1976) and 33.8 cm (1977) at the Sylvia Grinnell River (Table 7, Fig. 13). A high percentage of the catch from the Sylvia Grinnell River (93% in 1976, 84% in 1977) was retained compared with that at the Tree (24%) and Robertson (37%) rivers. Apparently anglers feel that if they are lucky enough to catch a charr at the Sylvia Grinnell River it is kept, whereas at the other locations mentioned, charr are plentiful and success is high, so most are released.

Compared with previous years, the size of charr angled at the Sylvia Grinnell River has declined drastically. Hunter (1958) reported 634 charr taken by anglers in 1958 (Table 7). Although he had no data on size of the angled catch he assumed it was similar to the mean size (2.21 kg) taken by the experimental gillnet fishery which took place in 1958. In 1976 census takers recorded a minimum of 212 charr kept from 229 landed during the period July 10 to August 19 (Table 7). The mean round weight per charr was 0.49 kg (N=137). In 1977, 379 of 415 charr landed during the period July 23 to August 15 were kept. The mean round weight per charr was 0.44 kg (N=259)(Table 7). According to Reiger (1982): "At one time eight-pound charr were abundant in the Sylvia Grinnell: last summer [1981] fish of any size were scarce."

## SUBSISTENCE FISHERY

No data were collected from the subsistence [domestic] gillnet fishery in 1976. Subsistence gillnetting occurred below the falls during the latter half of August, 1977, and samples of the catch were secured on August 23 and 25 (Fig. 14). The mean length of these charr was 38.9 cm (N = 53) and mean weight was 0.73 kg (N = 53).

Difficulties were experienced in determining daily catches. Fishermen were approached but some were reluctant to allow their catch to be counted. Counts were then made at a distance while the catch was being removed from the nets. Most nets were about 32 m long with 89 mm mesh size (stretched measure). A daily record of the number of gillnets set in the river was kept beginning August 17 when this activity started. Catches were heavy initially, when the upstream migration of charr began but tapered off near the end of the month (Table 8). About 2 282 charr were captured by the subsistence gillnet fishery.

Snagging began on August 23, 1976 and on August 16, 1977. Since snagging does not lure the fish to the hook but depends upon impaling it by chance, it is successful only when applied to concentrations of fish. Snagging success increased dramatically at the time of the spring tide (Fig. 11,12), indicating that the charr had concentrated in the river on their upstream migration. As discussed previously, the spring tide is necessary to enable the charr to surmount the falls on their way upstream. Hence the upstream migration of charr in the Sylvia Grinnell River is predictable.

At least 1 920 charr were snagged in 1976 and 4 923 in 1977. Mean fork length of snagged charr taken in 1976 was 22.5 cm (N = 644) and 32.8 cm (N = 292) in 1977. Mean round weight calculated from the weight-length relationship was 0.37 kg.

Charr taken by snagging were much smaller in 1976 than in 1977 (Fig. 15) although the reason for this is not clear. Generally it appears that small charr could be selectively taken by snagging since they are virtually powerless to escape once impaled on the hook. On the other hand, large charr could tear loose if superficially hooked. Some large charr taken by gillnets bore marks resembling those made by snag hooks. On August 21 and August 26, 1977, the snagging technique was carefully observed and 41% and 32%, respectively, of charr hooked escaped before they could be landed.

The most disturbing feature of the snagging fishery is the waste of charr. This was readily apparent to the observers. Many of the charr taken were considered too small to keep and were simply discarded on the rocks. A comparison of the size of charr taken by the different fisheries is shown in Fig. 14.

## SYLVIA GRINNELL LAKE

Most charr taken in Sylvia Grinnell Lake on August 8 and 9, 1977 were under 50 cm in length (Fig. 16). At this time the anadromous component of the stock would still be at sea. Hence this sample is probably comoosed of non-anadromous charr and perhaps some anadromous spawners.

## EFFECT OF FISHERIES ON THE CHARR STOCK

Hunter (1976) outlined the history of exploitation of this charr stock up to 1967. Fishing was limited to a traditional Inuit harvest until 1947, when commercial fishing began. Grainger's (1953) data indicate that the charr stock was relatively strong at the outset of commercial fishing. Grainger (1948) reported that a total of 7 700 charr were netted in 1948. Wright (1950) stated 2 167 charr were taken by the commercial fishery in 1950. Mean round weight per charr was 2.95 kg hence approximately 22 700 kg were taken in 1948 and 6 400 kg in 1950.

The commercial fishery did not operate from 1951 to 1957 but charr were taken by the subsistence and recreational fisheries (Hunter 1976). Hunter (1958) estimated the combined annual harvest from these two fisheries to be 12 000 kg in 1958. Commercial fishing was reestablished in 1958, this time by the Department of Northern Affairs, and an experimental gillnet fishery was conducted to determine the status of the stock (Hunter 1958). As shown in Fig. 3, the stock had not changed significantly since 1948. Apparently the first commercial fishery did not deplete the stock nor did the subsistence and recreational fisheries that followed immediately afterward.

On the strength of the 1958 evaluation an annual quota of 4 500 kg was allowed for the commercial fishery (Hunter 1976). It was estimated that the stock could sustain a combined commercial, recreational and subsistence harvest of 20 400 kg annually (Hunter 1958). Commercial production is shown in Table 9. Initially the quota was taken easily within a few days of the onset of fishing each year. However, catch per unit of effort declined steadily until commercial fishing was no longer worthwhile and the fishery ceased after 1966 (Hunter 1976). If the subsistence and recreational fisheries are assumed to have taken 12 000 kg per year during the operation of the commercial fishery, a total annual harvest of nearly 18 000 kg was likely at the outset. Apparently the charr stock could not sustain this yield and the fishery collapsed.

The subsistence and recreational fisheries that persisted after the cessation of commercial fishing still harvested large numbers of charr (Table 10) although the total landed weight was far less than previously taken, owing to the drastic reduction in the mean size of the charr. This continued fishing is probably responsible for the failure of the charr stock to recover to this day. Personnel from the Department of Fisheries and Oceans (DFO) observed this fishery during the summer of 1982 and results indicate that the stock had not yet recovered (W. Kowalczyk pers. comm.). A limited amount of gillnetting by DFO staff between August 19 and 31 (38, 63, 89 and 114 mm mesh size) captured charr with a mean length of 31.9 cm (N=32) and a mean age of 9 yr (N=27)(Fig. 17). Mean weight was 0.45 kg (N=31). These data were not used in the earlier comparison owing to the small sample size; however, they do support the observation that the stock is still in a reduced state.

Compared to other fishing activities, angling was responsible for harvesting the least number of charr although this activity was pursued over the longest period of time (Table 11). Four hundred and fifteen charr weighing a total of 181 kg were taken, with a mean round weight per charr of 0.44 kg. Subsistence gillnetting began with the first signs of the run near the spring tide in August and at least 2 282 charr weighing a total of 1 666 kg were taken. The mean weight per charr was 0.73 kg. Snagging was responsible for the harvest of the most charr. At least 4 923 were taken weighing a total of 1 822 kg. These charr were also the smallest with a mean weight per charr of only 0.37 kg. As a result many were discarded.

## RECOVERY OF THE CHARR STOCK

Charr grow very slowly, thus prerecruits to the fishable stock are composed of many year classes (Hunter 1976). Heavy exploitation of the fishable stock does not affect recruitment immediately. Recovery of the fishable stock can be quite rapid if fishing is ceased and the prerecruits are allowed to grow to fishable size. Such a rapid recovery took place with the commercial charr fishery at the Ekalluk River, Victoria Island (Kristofferson and Carder 1980).

There, fishing ceased in 1969 when the size of the charr had declined to a point where they were of little commercial value (Barlishen and Webber 1973). After a closure of three years, during which time the charr grew to an acceptable size, the fishery resumed in 1973 and remains viable today. However, the situation at the Sylvia Grinnell River could differ. Since the commercial fishery at the Ekalluk River is restricted to gillnets with a minimum mesh size of 139 mm, only the large charr are vulnerable to the gear. That is, the prerecruits were protected from the fishery and remained abundant. At the Sylvia Grinnell River even small charr are vulnerable to the gear, especially to snagging, and thus are recruited to the fishery at a very small size (Fig. 3) and early age (Fig. 4), well before the onset of sexual maturity. As a result, numbers of prerecruits are reduced, as is spawning potential. In effect, the recovery capacity of the charr stock in the Sylvia Grinnell River may be depressed. All that appears to remain of the robust stock Grainger (1953) observed in 1948/50/51 is the small, slow-growing mode of charr that was discussed in detail earlier. Whether these charr represent a distinct stock with a fixed, slow growth rate or whether individuals in this mode still have the capacity to rejuvenate the large mode through accellerated growth, if given the chance, is not known. In order to determine this, fishing pressure must be reduced.

The most effective way to determine if recovery is possible would be to cease all fishing activity at the river. Signs of recovery, if it does occur, should be evident within three to five years. However, such a closure would bring unnecessary hardships upon the community of Frobisher Bay. Angling is an important recreational pastime here and does not harvest large numbers of charr (Table 11), therefore allowing it to continue probably would not interfere significantly with the recovery of the stock. Subsistence fishing is, no doubt, very important to a small number of fishermen in Frobisher Bay, yet it is this fishery that is largely responsible for the prevention of the recovery of the stock. Ideally, if an alternate source of charr could be secured to fulfill this subsistence need, gillnetting could cease and recovery would be enhanced. An ample supply of charr is produced by the Pangnirtung commercial fishery 300 km away. Unless the destructive and wasteful practise of snagging is discontinued, hopes for recovery of this stock are bleak. Charr are extremely vulnerable when they congregate at the base of the falls on their upstream migration. Closure of this area of the river to all types of fishing is necessary to protect the stock. This is the only area where snagging is effective, and therefore, closure here will eliminate this practise. If incorporated, this strategy could contribute to the recovery of the stock. It is difficult to determine the time required to effect a recovery since it is not really known how the small size mode will respond to reduced fishing pressure. Any recovery will probably depend to a large extent on the continued intensity of fishing. If only angling is allowed, a measure of recovery could be effected in three to five years. Annual experimental gillnetting would provide a means to monitor the progress of the recovery. Once recovery is achieved assignment of a quota would be necessary to maintain the long-term viability of the fishery.

The future of this charr fishery depends upon whether or not this recommended program receives the support of the people of Frobisher Bay. A public meeting was held in Frobisher Bay on November 28, 1978, where DFO personnel explained the results of this study in detail to those present. Various alternatives were discussed, including total and partial closure of the river to fishing. In subsequent discussions it was decided that a questionnaire would be circulated by mail to the citizens of Frobisher Bay asking for their concerns and suggestions regarding the future management of the Sylvia Grinnell River. Response to the questionnaire was very poor. Another public meeting was held in Frobisher Bay on January 22, 1981. Once again DFO explained the results of this study and the alternatives for recovery were outlined. The Frobisher Bay Hunters and Trappers Association (HTA) was concerned with the small size of the charr in the river but did not express a willingness to support a closure of any kind. However, at an HTA meeting in Frobisher Bay held on June 10, 1982, and attended by DFO personnel, some of the people appeared to be amenable to a closure of the river to all methods of fishing from the falls to a point 100 m downstream. Recovery of the Sylvia Grinnell River charr stock to its former productive state will take place only with the cooperation of all parties involved in the fishery.

## RECOMMENDATIONS

1. It is recommended that the Sylvia Grinnell River be closed to all types of fishing from the head of the falls to a point not less than 100 m downstream.
2. It is recommended that an alternate source of Arctic charr be secured to satisfy the subsistence needs of the Inuit of Frobisher Bay, so that subsistence gillnet fishing can cease until the stock recovers. This will shorten the recovery period.
3. It is recommended that the minimum mesh size for subsistence gillnets be restricted to 139 mm, stretched measure, if recommendation #2 is not implemented. The annual catch from the subsistence gillnet fishery is to be monitored in this event.
4. It is recommended that an experimental gillnet program be carried out annually on the upstream migration of charr, in order to determine the recovery rate of the stock.

## ACKNOWLEDGMENTS

The authors wish to express their appreciation to Messrs. T. Chowns, E. Land, S. Ransom, D. Boxer and M. Ferguson of the Wildlife Service, Government of the Northwest Territories for their assistance in dealing with this situation. L. Dahlke, D. Leroux, D. Larsen, D. McGowan, G. Lacho and D. Rudy collected the data

for this program. S. Akeeagok, R. Allen, W. Kowalczyk, and G. Williams, DFO, Frobisher Bay, are thanked for their assistance. Ages of Arctic charr were determined by G. Carder. G. Yaremchuk assisted in the computer analyses. Drafting was done by Graphics Services, Freshwater Institute. Typing was done by S. Ahlgren. R. W. Moshenko and P. Richard reviewed an earlier draft of this report. Dr. R.A. Bodaly and W. Bond reviewed the final draft and provided many helpful suggestions.

## REFERENCES

BARLISHEN, W.J., and T.N. WEBBER. 1973. A history of the development of commercial fishing in the Cambridge Bay area of the Northwest Territories. Unpublished report for the Federal-Territorial Task Force on Fisheries Development in the Northwest Territories. 37 p.

DAHLKE, L.W., and M.R. FALK. 1979. Data on Arctic char from the Jayco River, Northwest Territories, 1975 . Can. Fish. Mar. Serv. Data Rep. 156: iv + 11 p.

de MARCH, L., B. de MARCH, and W. EDDY. 1978. Limnological, fisheries and stream benthic studies at Stanwell-Fletcher Lake, a large high Arctic lake. Can. Dep. Ind. North. Aff. ESCOM Report AI-04: xiv + 84 p.

FALK, M.R., and L.W. DAHLKE. 1979. Data on the Arctic char sport fishery at Tree River, Northwest Territories, 1978. Can. Fish. Mar. Serv. Data Rep. 153: iv + 32 p.

GRAINGER, E.H. 1948. Report on biological observations on Arctic char (Salvelinus alpinus) Frobisher Bay, 1948. Fish. Res. Board Can. Annual Report of Eastern Arctic Fisheries Investigations for 1948.

GRAINGER, E.H. 1953. On the age, growth, migration, reproductive potential and feeding habits of the Arctic char (Salvelinus alpinus) of Frobisher Bay, Baffin Island, Canada. J. Fish. Res. Board Can. 10: 326-370.

HUNTER, J.G. 1958. Production of Arctic char in the Sylvia Grinnell River, Baffin Island. Fish. Res. Board Can. Arctic Unit Annual Report and Investigators' Summaries, 1958.

HUNTER, J.G. 1963. Arctic char studies - Sylvia Grinnell River. Fish. Res. Board Can. Arctic Unit Annual Report and Investigators' Summaries April 1, 1962 to March 31, 1963.

HUNTER, J.G. 1976. Arctic char and hydroelectric power in the Sylvia Grinnell River. Fish. Res. Board Can. Manuscr. Rep. 1376: 21 p.

JOHNSON, L. 1976. Ecology of arctic populations of lake trout, Salvelinus namaycush, lake whitefish, Coregonus clupeaformis, Arctic charr, S. alpinus, and associated species in unexploited lakes in the Canadian Northwest Territories. J. Fish. Res. Board Can. 33: 2459-2488.

JOHNSON, L. 1980. The Arctic charr, Salvelinus alpinus, p. 15-98. In E.K. Balon (ed.) Charrs; salmonid fishes of the genus Salvelinus. Dr. W. Junk, The Hague.

KRISTOFFERSON, A.H., and G.W. CARDER. 1980. Data from the commercial fishery for Arctic char, Salvelinus alpinus (Linnaeus), in the Cambridge Bay area, Northwest Territories, 1971-78. Can. Data Rep. Fish. Aquat. Sci. 184: v + 25 p.

MOORE, J.W. 1975. Distribution, movements, and mortality of anadromous Arctic char, Salvelinus alpinus, in the Cumberland Sound area of Baffin Island. J. Fish Biol. 7: 339-348.

MOSHENKO, R.W. 1981. A preliminary assessment of the Arctic charr sport fishery on the Robertson River (Koluktoo Bay), Northwest Territories, 1979. Can. Data Rep. Fish. Aquat. Sci. 306: iv + 9 p.

MOSHENKO, R.W., R.F. PEET, L.W. DAHLKE, and D.H. DOWLER. 1983. The Arctic charr sport fishery at Tree River, Northwest Territories, Canada, 1964-78. In L. Johnson, R. McV. Clarke and K.E. Marshall (ed.) Biology of the Arctic charr, Proceedings of the International Symposium on Arctic charr, Winnipeg, Manitoba, May 1981. University of Manitoba Press, Winnipeg.

REIGER, G. 1982. River of record char. Fishing World 29(2): 82-94.

RICKER, W.E. 1977. The historical development, p. 1-26. In J.A. Gulland (ed.) Fish population dynamics. John Wiley and Sons, New York.

SCOTT, W.B., and E.J. CROSSMAN. 1973. Freshwater fishes of Canada. Fish. Res. Board Can. Bull. 184: xiii + 955 p.

SEKERAK, A.D., D. THOMSON, H. BAIN and J. ACREMAN. 1976. Summer surveys of the marine ecology of Creswell Bay, Somerset Island and Assistance Bay, Cornwallis Island, N.W.T. 1975. LGL Ltd. environmental research associates, Toronto, Ontario. 215 p.

SPRULES, W.M. 1952. The Arctic char of the west coast of Hudson Bay. J. Fish. Res. Board Can. 9: 1-15.

STATISTICAL ANALYSIS SYSTEM. 1979. SAS user's guide. 1979 edition. Cary, North Carolina.

TYLER, A.V., and V.F. GALLUCCI. 1980. Dynamics of fished stocks, p. 111-147. In R.T. Lackey and L.A. Nielsen (ed.) Fisheries Management. John Wiley and Sons, New York.

WRIGHT, J. 1950. Arctic char, Frobisher Bay and Atlantic cod, Resolution Island. Fish. Res. Board Canada Annual Report of Eastern Arctic Fisheries Investigations for 1950.




Fig. 1. Map of the Frobisher Bay area, Baffin Island, showing the Sylvia Grinnell River.




Fig. 2. Map of the outlet of the Sylvia Grinnell River showing the study area.




Fig. 3. A comparison of length-frequency distributions of samples of Arctic charr taken from the Sylvia Grinnell River during 1948/50/51, 1958 and 1976/77.




Fig. 4. A comparison of age-frequency distributions of samples of Arctic charr taken from the Sylvia Grinnell River during 1948/50/51 and 1976/77.




Fig. 5. Catch curves for Arctic charr taken from the Sylvia Grinnell River during 1948/50/51, 1958 and 1976/77.




Fig. 6. A comparison of mortality rates (catch curve) for Arctic charr, aged 13(14) to 18 years, taken from the Sylvia Grinnell River during 1948/50/51, 1958 and 1976/77.




Fig. 7. A comparison of growth rates of Arctic charr taken from the Sylvia Grinnell River in 1948/50/51 and 1976/77 with those of Arctic charr from other rivers in the Northwest Territories.




Fig. 8. A comparison of length- and age-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by angling in 1976 and 1977.




Fig. 9. A comparison of length- and age-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by experimental gillnet in 1976 and 1977.




Fig. 10. Length-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by experimental gillnets on four occasions during 1977.




Fig. 11. Daily catch of Arctic charr from the Sylvia Grinnell River taken by angling and snagging compared with daily high tide level, 1976.




Fig. 12. Daily catch of Arctic charr from the Sylvia Grinnell River taken by angling and snagging compared with daily high tide level, 1977.




Fig. 13. A comparison of length-frequency distributions of Arctic charr taken by angling from the Sylvia Grinnell River in 1976/77, the Tree River in 1978 and the Robertson River in 1979.




Fig. 14. A comparison of length-frequency distributions of Arctic charr taken by different types of gear from the Sylvia Grinnell River, 1976/77.




Fig. 15. A comparison of length-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by snagging in 1976 and 1977.




Fig. 16. Length-frequency distribution of Arctic charr taken from Sylvia Grinnell Lake by experimental gillnet in 1977.




Fig. 17. Length- and age-frequency distributions of Arctic charr taken from the Sylvia Grinnell River by experimental gillnet in 1982.

Table 1. Instantaneous total and fishing mortality and annual survival of Arctic charr, aged 13-18 years, from the Sylvia Grinnell River in 1948/50/51, 1958 and 1976/77.

| Year(s) | Age Groups (years) | Instantaneous Total Mortality (Catch Curve) Z | Instantaneous Fishing Mortality (Z-0.13) F | Annual Survival S(%) |
|---|---|---|---|---|
| 1948/50/51 | 13-18 | 0.31 | 0.18 | 73 |
| 1958 | 14-18 | 0.14 | 0.01[a] | 87 |
| 1976/77 | 13-18 | 0.70 | 0.57 | 50 |

[a]Sport and subsistence fishing is known to have taken place prior to and during 1958 so fishing mortality did occur. However, it was probably light. The estimation of instantaneous natural mortality (0.13) for these age groups may be too high.

Table 2. Weight-length relationships of Arctic charr from a number of locations compared with those from the Sylvia Grinnell River, 1976 and 1977. Log W = log a + n log L, where W=weight in grams, L=fork length in millimetres. Data tabulated by Johnson (1980).

| Location | Date | n | log a |
|---|---|---|---|
| Sylvia Grinnell R. | 10/7 to 28/8/76 | 3.0724 | -5.2101 |
| Sylvia Grinnell R. | 23/7 to 28/8/77 | 3.1368 | -5.3254 |
| Robertson R.[a] | 26/8/73 | 3.0041 | -4.9933 |
| Creswell Bay[b] | 30/7 to 13/8/75 | 3.2653 | -5.6875 |
| Nauyuk Lake[c] | 15/6 to 15/7/75 | 3.1668 | -5.5062 |
| Nauyuk Lake[c] | 1/8 to 10/9/75 | 3.1077 | -5.2281 |

[a]Peet (unpublished).

[b]Sekerak et al. (1976).

[c]Johnson (unpublished).

Table 3. Condition factor (K) of Arctic charr caught in the Sylvia Grinnell River in 1976 and 1977.

| Length Interval | 1976 | | 1977 | |
|---|---|---|---|---|
| mm | N | K | N | K |
| 150 - 199 | 20 | 0.89 | 9 | 1.03 |
| 200 - 249 | 39 | 0.95 | 34 | 1.03 |
| 250 - 299 | 24 | 0.96 | 124 | 1.01 |
| 300 - 349 | 69 | 0.91 | 325 | 1.02 |
| 350 - 399 | 45 | 0.97 | 190 | 1.10 |
| 400 - 449 | 41 | 0.95 | 98 | 1.11 |
| 450 - 499 | 15 | 0.98 | 50 | 1.13 |
| 500 - 549 | 2 | 1.08 | 16 | 1.17 |
| 550 - 599 | - | - | 1 | 1.12 |
| TOTAL | 255 | | 847 | |
| MEAN | | 0.94 | | 1.06 |

Table 4. Age, mean length and size range of Arctic charr from the Sylvia Grinnell River, 1948, 1950 and 1951. From Grainger (1953).

| Age (yr) | Number | Mean Length (cm) | Length Range (cm) |
|---|---|---|---|
| 4 | 3 | 9.0 | 8.5 - 10.0 |
| 5 | 18 | 13.0 | 11.0 - 14.5 |
| 6 | 7 | 13.9 | 12.5 - 16.5 |
| 7 | 7 | 17.2 | 14.5 - 28.5 |
| 8 | 25 | 30.2 | 17.0 - 39.5 |
| 9 | 68 | 33.7 | 18.5 - 46.5 |
| 10 | 68 | 36.3 | 26.5 - 52.0 |
| 11 | 73 | 41.4 | 29.5 - 55.0 |
| 12 | 85 | 42.5 | 30.5 - 57.0 |
| 13 | 96 | 48.4 | 30.5 - 65.0 |
| 14 | 71 | 51.0 | 36.0 - 63.0 |
| 15 | 48 | 55.3 | 31.5 - 67.5 |
| 16 | 38 | 55.7 | 44.0 - 68.5 |
| 17 | 31 | 55.9 | 36.5 - 70.0 |
| 18 | 19 | 60.0 | 46.0 - 70.5 |
| 19 | 23 | 61.3 | 46.5 - 74.0 |
| 20 | 35 | 64.9 | 55.0 - 76.5 |
| 21 | 35 | 64.2 | 50.0 - 76.5 |
| 22 | 19 | 65.8 | 44.0 - 78.5 |
| 23 | 16 | 67.8 | 51.0 - 81.5 |
| 24 | 9 | 67.8 | 58.5 - 78.5 |
| 24 plus | 12 | 68.5 | 51.0 - 82.5 |

Table 5. Percent of Arctic charr, by length interval, captured in spawning condition while migrating up the Sylvia Grinnell River in 1976/77.

| Length Interval (mm) | Males | | Females | |
|---|---|---|---|---|
| | N | % Mature | N | % Mature |
| 150 - 199 | 8 | 0 | 15 | 0 |
| 200 - 249 | 23 | 0 | 32 | 0 |
| 250 - 299 | 47 | 0 | 71 | 0 |
| 300 - 349 | 158 | 0 | 130 | 0 |
| 350 - 399 | 97 | 0 | 87 | 2 |
| 400 - 449 | 73 | 0 | 52 | 25 |
| 450 - 499 | 28 | 0 | 31 | 45 |
| 500 - 549 | 9 | 33 | 6 | 67 |
| 500 - 599 | 1 | 0 | - | - |
| TOTAL | 441 | 1 | 424 | 8 |

Table 6. Percent of Arctic charr, by age, captured in spawning condition while migrating up the Sylvia Grinnell River in 1976/77.

| Age (yr) | Males | | Females | |
|---|---|---|---|---|
| | N | % Mature | N | % Mature |
| 5 | 3 | 0 | - | - |
| 6 | 5 | 0 | 11 | 0 |
| 7 | 14 | 0 | 24 | 0 |
| 8 | 32 | 0 | 32 | 0 |
| 9 | 56 | 0 | 52 | 0 |
| 10 | 68 | 0 | 72 | 4 |
| 11 | 72 | 1 | 60 | 7 |
| 12 | 63 | 0 | 8 | 13 |
| 13 | 41 | 0 | 39 | 8 |
| 14 | 17 | 0 | 21 | 19 |
| 15 | 11 | 9 | 11 | 18 |
| 16 | 7 | 0 | 6 | 33 |
| 17 | 4 | 0 | 2 | 0 |
| 18 | 2 | 50 | - | - |
| 19 | - | - | - | - |
| 20 | - | - | 2 | 100 |
| TOTAL | 395 | >1 | 340 | 6 |

Table 7. A comparison of creel census data for Arctic charr angled in the Sylvia Grinnel River in 1958, 1976-77, the Tree River in 1978 and the Robertson River in 1979.

| Location | No. Anglers Censused | Total Catch | % Anglers Successful | No. Fish Retained (%) | Total Hours Fished | Catch/ Angler Hour | No. Fish Sampled | Mean Round Weight (kg) | Mean Fork Length (cm) |
|---|---|---|---|---|---|---|---|---|---|
| Sylvia Grinnell 1977 | 374 | 415 | 47 | 379 (84) | 1 102 | 0.4 | 259 | 0.44 | 33.8 |
| Sylvia Grinnell 1976 | 427 | 229 | 26 | 212 (93) | 871 | 0.3 | 137 | 0.49 | 36.4 |
| Sylvia Grinnell 1958 | 87 | 634 | - | - | 2 330 | 0.3 | 902[c] | 2.21 | 48.0 |
| Tree River[a] 1978 | 472 | 2 186 | 84 | 524 (24) | 4 495 | 0.5 | 573 | 4.90 | 75.6 |
| Robertson River[b] 1979 | 77 | 368 | 95 | 135 (37) | 307 | 1.2 | 19 | 3.00 | 65.9 |

[a] from Falk and Dahlke 1979.

[b] from Moshenko 1980.

[c] samples taken by experimental gillnet. See text for explanation.

Table 8. Catch record of the subsistence gillnet fishery in the Sylvia Grinnell River, August 1977.

| Date | No. Nets (ca 32m in length) | Hours Set | Catch | | Total Daily Catch |
|---|---|---|---|---|---|
| | | | Calculated | Counted | |
| Aug. 17 | 4 | 8 | 203[a] | | 203 |
| 18 | 5 | 8 | 253[a] | | 253 |
| 19 | 5 | 7 | 222[a] | | 222 |
| 20 | 7 | 8 | 355[a] | | 355 |
| 21 | 7 | 8 | 355[a] | | 355 |
| 23 | 1 | 10 | | 150 | 150 |
| 23 | 2 | 7 | 200[b] | 200 | 400 |
| 24 | 1 | 12 | | 130 | 130 |
| 24 | 1 | 6 | | 50 | 50 |
| 25 | 1 | 4 | | 16 | 16 |
| 25 | 1 | 4 | 15[c] | | 15 |
| 25 | 1 | 3 | | 10 | 10 |
| 25 | 1 | 24 | 89[c] | | 89 |
| 25 | 1 | 1 | 4[c] | | 4 |
| 25 | 1 | 2.5 | 9[c] | | 9 |
| 26 | 1 | 1 | | 15 | 15 |
| 26 | 1 | 24 | | 3 | 3 |
| 27 | 1 | 24 | 3[d] | | 3 |
| 28 | 5 | ——— No Observations Made ——— | | | |
| TOTAL | | | | | 2 282 |

[a]Calculated using catch per unit of effort of 89mm panel from experimental gillnet set of August 15.

[b]Assumed catch similar to that counted from adjacent net.

[c]Calculated using catch per unit of effort of other gillnets lifted August 25 where catch was counted.

[d]Assumed catch similar to that of August 26.

Table 9. Annual harvest statistics for Arctic charr from the Sylvia Grinnell River, 1958-66.

| Year | Commercial[a] Harvest kg | Total Harvest (Includes estimated subsistence and recreational harvest of 12,000 kg)[b] kg |
|---|---|---|
| 1958 | 5 834 | 17 834 |
| 1959 | 9 803 | 21 803 |
| 1960 | 5 532 | 17 532 |
| 1961 | 4 674 | 16 674 |
| 1962 | 4 688 | 16 688 |
| 1963 | 4 920 | 16 920 |
| 1964 | 3 824 | 15 824 |
| 1965 | 5 588 | 17 588 |
| 1966 | 4 658 | 16 658 |

[a]From Hunter (1958) and Hunter (1976). Quota was 4 500 kg.

[b]From Hunter (1958) for 1958. It is assumed that this level of harvest continued until 1966.

Table 10. Annual harvest, mean round weight per charr and number of charr taken by fisheries at the Sylvia Grinnell River in 1948, 1950, 1958 and 1977.

| Year | Annual Harvest (kg) | Mean Round Weight per Charr (kg) | Number of Charr |
|---|---|---|---|
| 1948[a] | 22 700 | 2.95 | 7 700 |
| 1950[b] | 6 400 | 2.95 | 2 167 |
| 1958[c] | 17 834 | 2.20 | 8 106 |
| 1977 | 3 667 | 0.48 | 7 620 |

[a]From Grainger (1948).

[b]From Wright (1950).

[c]From Hunter (1958).

Table 11. Number of days fished, harvest, mean round weight per charr and number of charr taken by angling, subsistence gillnetting and snagging at the Sylvia Grinnell River, 1977.

| Fishery | No. Days Fished | Harvest (kg) | Mean Round Weight per Charr (kg) | No. Charr |
|---|---|---|---|---|
| Angling | 22 | 181 | 0.44 | 415 |
| Subsistence Gillnets | 11 | 1 666 | 0.73 | 2 282 |
| Snagging | 13 | 1 822 | 0.37 | 4 923 |
| TOTAL | | 3 669 | | 7 620 |

Appendix 1. Summary of biological data by length interval for Arctic charr taken by angling in the Sylvia Grinnell River, 1976.

| LENGTH INTERVAL (MM) | MALES | | | | | | FEMALES | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | | % |
| | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT |
| 150 | - | - | - | - | - | - | 1 | 199 | 75 | - | 0.95 | 0 | 1 | 199 | 75 | - | 0.95 | 0 |
| 200 | - | - | - | - | - | - | 1 | 245 | 100 | - | 0.68 | 0 | 1 | 245 | 100 | - | 0.68 | 0 |
| 250 | 7 | 275 | 186 | 56 | 0.87 | 0 | 7 | 285 | 221 | 37 | 0.96 | 0 | 14 | 280 | 204 | 49 | 0.92 | 0 |
| 300 | 27 | 324 | 316 | 65 | 0.92 | 0 | 24 | 326 | 302 | 54 | 0.87 | 0 | 51 | 325 | 309 | 60 | 0.90 | 0 |
| 350 | 15 | 367 | 462 | 88 | 0.93 | 0 | 12 | 378 | 496 | 69 | 0.92 | 0 | 27 | 371 | 477 | 81 | 0.92 | 0 |
| 400 | 17 | 424 | 682 | 107 | 0.89 | 0 | 11 | 420 | 711 | 111 | 0.96 | 9 | 28 | 422 | 694 | 108 | 0.92 | 4 |
| 450 | 5 | 464 | 955 | 148 | 0.95 | 0 | 9 | 472 | 1017 | 185 | 0.96 | 11 | 14 | 470 | 995 | 170 | 0.96 | 7 |
| 500 | 1 | 536 | 1625 | - | 1.06 | 0 | - | - | - | - | - | - | 1 | 536 | 1625 | - | 1.06 | 0 |
| TOTAL | 72 | | | | | | 65 | | | | | | 137 | | | | | |
| MEAN | | 364 | 483 | 265 | 0.91 | | | 364 | 491 | 286 | 0.91 | | | 364 | 487 | 274 | 0.91 | |



Appendix 2. Summary of biological data by age for Arctic charr taken by angling in the Sylvia Grinnell River, 1976.

| AGE (YR) | MALES | | | | | | | FEMALES | | | | | | | COMBINED | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % |
| | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT |
| 7 | 1 | 324 | - | 250 | - | 0.74 | 0 | 1 | 199 | - | 75 | - | 0.95 | 0 | 2 | 262 | 88.4 | 163 | 124 | 0.84 | 0 |
| 8 | 6 | 310 | 24.9 | 283 | 68 | 0.94 | 0 | 1 | 265 | - | 200 | - | 1.07 | 0 | 7 | 304 | 28.4 | 271 | 70 | 0.96 | 0 |
| 9 | 3 | 307 | 4.0 | 250 | 50 | 0.86 | 0 | 6 | 289 | 29.9 | 229 | 87 | 0.91 | 0 | 9 | 295 | 25.5 | 236 | 74 | 0.89 | 0 |
| 10 | 10 | 334 | 60.4 | 380 | 237 | 0.90 | 0 | 12 | 346 | 45.2 | 419 | 211 | 0.94 | 0 | 22 | 340 | 51.7 | 401 | 219 | 0.92 | 0 |
| 11 | 12 | 357 | 61.6 | 452 | 264 | 0.92 | 0 | 13 | 338 | 28.0 | 337 | 116 | 0.84 | 0 | 25 | 347 | 47.1 | 392 | 205 | 0.88 | 0 |
| 12 | 15 | 364 | 44.0 | 465 | 207 | 0.92 | 0 | 8 | 361 | 46.4 | 447 | 197 | 0.90 | 0 | 23 | 363 | 43.8 | 459 | 199 | 0.91 | 0 |
| 13 | 8 | 399 | 47.3 | 619 | 206 | 0.94 | 0 | 5 | 403 | 64.1 | 635 | 327 | 0.90 | 20 | 13 | 401 | 51.8 | 625 | 246 | 0.92 | 8 |
| 14 | 5 | 403 | 48.3 | 610 | 204 | 0.91 | 0 | 3 | 435 | 54.5 | 833 | 325 | 0.98 | 0 | 8 | 415 | 49.6 | 694 | 260 | 0.93 | 0 |
| 15 | 4 | 446 | 77.7 | 919 | 515 | 0.95 | 0 | 2 | 421 | 22.6 | 650 | 141 | 0.86 | 0 | 6 | 438 | 62.3 | 829 | 427 | 0.92 | 0 |
| 16 | 2 | 413 | 10.6 | 600 | 71 | 0.85 | 0 | 2 | 475 | 12.7 | 1200 | 71 | 1.12 | 0 | 4 | 444 | 37.3 | 900 | 351 | 0.99 | 0 |
| 17 | 2 | 407 | 7.1 | 550 | 71 | 0.81 | 0 | 2 | 472 | 21.9 | 1075 | 35 | 1.03 | 0 | 4 | 439 | 39.5 | 813 | 307 | 0.92 | 0 |
| TOTAL | 68 | | | | | | | 55 | | | | | | | 123 | | | | | | |
| MEAN | | 365 | 60 | 483 | 267 | 0.91 | | | 358 | 65.8 | 475 | 300 | 0.92 | | | 362 | 62.3 | 480 | 281 | 0.91 | |
| MEAN AGE | 11.6 | | | | | | | | | | | | | | | | | | | | |

Appendix 3. Summary of biological data by length interval for Arctic charr taken by angling in the Sylvia Grinnel River, 1977.

| LENGTH INTERVAL (MM) | MALES | | | | | | FEMALES | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | | % |
| | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT |
| 100 | - | - | - | - | - | - | 1 | 127 | 150 | - | 7.32 | 0 | 1 | 127 | 150 | - | 7.32 | 0 |
| 150 | - | - | - | - | - | - | - | - | - | - | - | - | 1 | 181 | 50 | - | 0.84 | 0 |
| 200 | 3 | 223 | 208 | 88 | 1.85 | 0 | 1 | 246 | 125 | - | 0.84 | 0 | 6 | 231 | 167 | 72 | 1.39 | 0 |
| 250 | 24 | 285 | 225 | 31 | 0.97 | 0 | 43 | 282 | 222 | 37 | 0.99 | 0 | 67 | 283 | 223 | 35 | 0.98 | 0 |
| 300 | 49 | 324 | 336 | 63 | 0.98 | 0 | 48 | 324 | 334 | 70 | 0.97 | 0 | 98 | 324 | 334 | 66 | 0.97 | 0 |
| 350 | 17 | 370 | 516 | 84 | 1.01 | 0 | 24 | 370 | 540 | 95 | 1.06 | 0 | 42 | 371 | 532 | 90 | 1.04 | 0 |
| 400 | 16 | 424 | 802 | 155 | 1.04 | 0 | 13 | 427 | 860 | 81 | 1.10 | 38 | 29 | 425 | 828 | 128 | 1.07 | 17 |
| 450 | 5 | 476 | 1080 | 127 | 1.00 | 0 | 6 | 465 | 1058 | 153 | 1.05 | 50 | 11 | 470 | 1068 | 135 | 1.02 | 27 |
| 500 | 2 | 523 | 1575 | 35 | 1.10 | 0 | 1 | 500 | 1300 | - | 1.04 | 0 | 3 | 515 | 1483 | 161 | 1.08 | 0 |
| 550 | 1 | 570 | 2075 | - | 1.12 | 0 | - | - | - | - | - | - | 1 | 570 | 2075 | - | 1.12 | 0 |
| TOTAL | 117 | | | | | | 137 | | | | | | 259 | | | | | |
| MEAN | | 346 | 468 | 321 | 1.02 | | | 334 | 421 | 257 | 1.05 | | | 338 | 438 | 288 | 1.03 | |



Appendix 4. Summary of biological data by age for Arctic charr taken by angling in the Sylvia Grinnell River, 1977.

| AGE (YR) | MALES | | | | | | | FEMALES | | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % |
| | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT |
| 5 | 1 | 281 | - | 200 | - | 0.90 | 0 | - | - | - | - | - | - | - | 1 | 281 | - | 200 | - | 0.90 | 0 |
| 6 | 1 | 223 | - | 125 | - | 1.13 | 0 | 2 | 267 | 29.7 | 163 | 53 | 0.84 | 0 | 3 | 252 | 33.0 | 150 | 43 | 0.93 | 0 |
| 7 | 2 | 246 | 51.6 | 213 | 18 | 1.60 | 0 | 3 | 271 | 18.9 | 242 | 52 | 1.20 | 0 | 5 | 261 | 32.3 | 230 | 41 | 1.36 | 0 |
| 8 | 11 | 304 | 26.3 | 300 | 90 | 1.04 | 0 | 13 | 298 | 32.0 | 277 | 119 | 1.00 | 0 | 26 | 294 | 38.7 | 272 | 115 | 1.01 | 0 |
| 9 | 18 | 318 | 37.9 | 342 | 177 | 0.99 | 0 | 21 | 314 | 34.9 | 321 | 130 | 0.99 | 0 | 39 | 316 | 35.9 | 331 | 152 | 0.99 | 0 |
| 10 | 14 | 344 | 61.5 | 475 | 360 | 1.03 | 0 | 23 | 326 | 46.9 | 374 | 204 | 0.99 | 4 | 37 | 332 | 52.8 | 412 | 273 | 1.01 | 3 |
| 11 | 22 | 319 | 40.5 | 342 | 174 | 1.02 | 0 | 22 | 335 | 48.1 | 406 | 217 | 1.00 | 9 | 45 | 325 | 46.2 | 368 | 199 | 1.01 | 4 |
| 12 | 15 | 368 | 50.1 | 542 | 249 | 1.02 | 0 | 9 | 337 | 45.0 | 403 | 239 | 0.98 | 0 | 24 | 356 | 49.7 | 490 | 249 | 1.01 | 0 |
| 13 | 9 | 409 | 80.0 | 781 | 562 | 0.99 | 0 | 14 | 373 | 61.0 | 588 | 319 | 1.04 | 7 | 23 | 387 | 69.6 | 663 | 429 | 1.02 | 4 |
| 14 | 4 | 383 | 35.7 | 519 | 206 | 0.88 | 0 | 5 | 351 | 48.9 | 475 | 267 | 1.00 | 0 | 9 | 365 | 44.2 | 494 | 228 | 0.95 | 0 |
| 15 | 3 | 454 | 55.2 | 925 | 307 | 0.96 | 0 | 2 | 374 | 5.7 | 513 | 18 | 0.98 | 0 | 5 | 422 | 58.9 | 760 | 314 | 0.97 | 0 |
| 16 | 1 | 395 | - | 550 | - | 0.89 | 0 | - | - | - | - | - | - | - | 1 | 395 | - | 550 | - | 0.89 | 0 |
| TOTAL | 101 | | | | | | | 114 | | | | | | | 218 | | | | | | |
| MEAN | | 340 | 62 | 445 | 308 | 1.02 | | | 328 | 49.7 | 388 | 222 | 1.00 | | | 332 | 57.4 | 410 | 267 | 1.01 | |
| MEAN AGE | 10.5 | | | | | | | | | | | | | | | | | | | | |

Appendix 5. Summary of biological data by length interval for Arctic charr taken by experimental gillnets (38, 63, 89, 114, and 139 mm mesh size) in the Sylvia Grinnell River, 1976.

| LENGTH INTERVAL (MM) | MALES | | | | | | FEMALES | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | N | LENGTH(MM) MEAN | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | WEIGHT(G) MEAN | SD | K | % MAT |
| 150 | 7 | 190 | 61 | 13 | 0.88 | 0 | 11 | 191 | 61 | 13 | 0.87 | 0 | 19 | 191 | 62 | 13 | 0.89 | 0 |
| 200 | 16 | 221 | 106 | 28 | 0.96 | 0 | 21 | 223 | 105 | 19 | 0.94 | 0 | 38 | 222 | 105 | 23 | 0.95 | 0 |
| 250 | 3 | 294 | 267 | 29 | 1.05 | 0 | 7 | 290 | 250 | 29 | 1.02 | 0 | 10 | 292 | 255 | 28 | 1.03 | 0 |
| 300 | 11 | 315 | 291 | 38 | 0.93 | 0 | 6 | 333 | 392 | 58 | 1.05 | 0 | 18 | 321 | 325 | 65 | 0.97 | 0 |
| 350 | 11 | 376 | 550 | 81 | 1.03 | 0 | 7 | 375 | 564 | 75 | 1.07 | 0 | 18 | 375 | 556 | 76 | 1.05 | 0 |
| 400 | 6 | 417 | 775 | 52 | 1.07 | 0 | 6 | 426 | 767 | 93 | 0.99 | 0 | 13 | 421 | 773 | 70 | 1.03 | 0 |
| 450 | 1 | 453 | 1150 | - | 1.24 | 0 | - | - | - | - | - | - | 1 | 453 | 1150 | - | 1.24 | 0 |
| 500 | - | - | - | - | - | - | 1 | 503 | 1400 | - | 1.10 | 0 | 1 | 503 | 1400 | - | 1.10 | 0 |
| TOTAL | 55 | | | | | | 59 | | | | | | 118 | | | | | |
| MEAN | | 296 | 327 | 267 | 0.98 | | | 280 | 287 | 282 | 0.97 | | | 288 | 307 | 275 | 0.98 | |

Appendix 6. Summary of biological data by age for Arctic charr taken by experimental gillnets in the Sylvia Grinnell River, 1976.

| AGE (YR) | MALES | | | | | | | FEMALES | | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | N | LENGTH(MM) MEAN | SD | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | SD | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | SD | WEIGHT(G) MEAN | SD | K | % MAT |
| 5 | 2 | 210 | 38.2 | 100 | 71 | 0.97 | 0 | - | - | - | - | - | - | - | 2 | 210 | 38.2 | 100 | 71 | 0.97 | 0 |
| 6 | 2 | 198 | 1.4 | 63 | 18 | 0.80 | 0 | 4 | 210 | 20.1 | 88 | 32 | 0.91 | 0 | 7 | 204 | 15.8 | 79 | 27 | 0.90 | 0 |
| 7 | 6 | 201 | 15.7 | 71 | 19 | 0.86 | 0 | 13 | 208 | 17.5 | 87 | 26 | 0.95 | 0 | 20 | 206 | 16.3 | 83 | 24 | 0.93 | 0 |
| 8 | 6 | 279 | 82.8 | 267 | 241 | 0.97 | 0 | 8 | 243 | 41.9 | 153 | 95 | 0.96 | 0 | 14 | 258 | 62.7 | 202 | 175 | 0.97 | 0 |
| 9 | 13 | 275 | 62.8 | 246 | 193 | 1.02 | 0 | 9 | 282 | 58.8 | 269 | 168 | 1.04 | 0 | 22 | 278 | 59.9 | 256 | 179 | 1.03 | 0 |
| 10 | 6 | 369 | 52.3 | 558 | 248 | 1.03 | 0 | 6 | 343 | 71.4 | 433 | 268 | 0.93 | 0 | 12 | 356 | 61.2 | 496 | 254 | 0.98 | 0 |
| 11 | 5 | 337 | 17.5 | 360 | 82 | 0.93 | 0 | 5 | 309 | 54.9 | 340 | 213 | 1.01 | 0 | 11 | 323 | 39.0 | 345 | 146 | 0.96 | 0 |
| 12 | 1 | 363 | - | 500 | - | 1.05 | 0 | 1 | 362 | - | 450 | - | 0.95 | 0 | 3 | 382 | 33.8 | 583 | 189 | 1.02 | 0 |
| 13 | 4 | 411 | 38.0 | 775 | 287 | 1.08 | 0 | 5 | 426 | 49.5 | 820 | 340 | 1.02 | 0 | 9 | 419 | 42.8 | 800 | 299 | 1.05 | 0 |
| 14 | 1 | 388 | - | 600 | - | 1.03 | 0 | 1 | 436 | - | 800 | - | 0.97 | 0 | 2 | 412 | 33.9 | 700 | 141 | 1.00 | 0 |
| 15 | - | - | - | - | - | - | - | 1 | 424 | - | 750 | - | 0.98 | 0 | 1 | 424 | - | 750 | - | 0.98 | 0 |
| 16 | 1 | 397 | - | 600 | - | 0.96 | 0 | - | - | - | - | - | - | - | 1 | 397 | - | 600 | - | 0.96 | 0 |
| TOTAL | 47 | | | | | | | 53 | | | | | | | 104 | | | | | | |
| MEAN | | 297 | 83 | 330 | 271 | 0.98 | | | 283 | 85.1 | 293 | 284 | 0.97 | | | 289 | 84.2 | 311 | 278 | 0.98 | |
| MEAN AGE | 9.2 | | | | | | | | | | | | | | | | | | | | |

Appendix 7. Summary of biological data by length interval for Arctic charr taken by experimental gillnets in the Sylvia Grinnell River, 1977.

| LENGTH INTERVAL (MM) | MALES | | | | | | FEMALES | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | N | LENGTH(MM) MEAN | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | WEIGHT(G) MEAN | SD | K | % MAT |
| 150 | 1 | 186 | 75 | - | 1.17 | 0 | 2 | 183 | 75 | 0 | 1.22 | 0 | 3 | 184 | 75 | 0 | 1.20 | 0 |
| 200 | 2 | 230 | 100 | 0 | 0.82 | 0 | 7 | 223 | 93 | 19 | 0.84 | 0 | 10 | 224 | 95 | 16 | 0.84 | 0 |
| 250 | 7 | 282 | 251 | 60 | 1.11 | 0 | 4 | 290 | 275 | 35 | 1.13 | 0 | 11 | 285 | 260 | 52 | 1.12 | 0 |
| 300 | 36 | 328 | 374 | 58 | 1.05 | 0 | 38 | 322 | 376 | 75 | 1.12 | 0 | 74 | 325 | 375 | 67 | 1.09 | 0 |
| 350 | 37 | 370 | 600 | 121 | 1.17 | 0 | 28 | 377 | 618 | 97 | 1.15 | 7 | 65 | 373 | 608 | 111 | 1.16 | 3 |
| 400 | 27 | 420 | 865 | 120 | 1.17 | 0 | 18 | 421 | 860 | 133 | 1.15 | 22 | 45 | 421 | 863 | 124 | 1.16 | 9 |
| 450 | 13 | 475 | 1306 | 85 | 1.22 | 0 | 15 | 468 | 1212 | 209 | 1.17 | 60 | 28 | 472 | 1255 | 168 | 1.19 | 32 |
| 500 | 5 | 531 | 1775 | 87 | 1.19 | 40 | 3 | 514 | 1692 | 577 | 1.22 | 100 | 8 | 525 | 1744 | 318 | 1.20 | 63 |
| TOTAL | 128 | | | | | | 115 | | | | | | 244 | | | | | |
| MEAN | | 377 | 679 | 388 | 1.13 | | | 365 | 628 | 387 | 1.13 | | | 371 | 652 | 388 | 1.13 | |



Appendix 8. Summary of biological data by age for Arctic charr taken by experimental gillnets in the Sylvia Grinnell River, 1977.

| AGE (YR) | MALES | | | | | | | FEMALES | | | | | | | COMBINED | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | N | LENGTH(MM) MEAN | SD | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | SD | WEIGHT(G) MEAN | SD | K | % MAT | N | LENGTH(MM) MEAN | SD | WEIGHT(G) MEAN | SD | K | % MAT |
| 6 | 2 | 278 | 130.1 | 350 | 389 | 1.20 | 0 | 3 | 195 | 21.4 | 83 | 14 | 1.13 | 0 | 5 | 228 | 80.7 | 190 | 243 | 1.16 | 0 |
| 7 | 1 | 297 | - | 350 | - | 1.34 | 0 | 3 | 253 | 29.3 | 175 | 109 | 1.00 | 0 | 5 | 256 | 33.6 | 195 | 120 | 1.05 | 0 |
| 8 | 4 | 316 | 64.3 | 463 | 309 | 1.24 | 0 | 8 | 278 | 52.0 | 244 | 136 | 0.99 | 0 | 12 | 291 | 56.5 | 317 | 222 | 1.07 | 0 |
| 9 | 10 | 341 | 50.5 | 450 | 248 | 1.04 | 0 | 7 | 329 | 35.9 | 418 | 162 | 1.12 | 0 | 17 | 336 | 44.2 | 437 | 211 | 1.08 | 0 |
| 10 | 28 | 372 | 60.1 | 658 | 320 | 1.16 | 0 | 21 | 354 | 36.9 | 537 | 191 | 1.16 | 5 | 49 | 364 | 51.8 | 606 | 276 | 1.16 | 2 |
| 11 | 19 | 365 | 53.5 | 601 | 337 | 1.14 | 0 | 16 | 363 | 57.6 | 589 | 336 | 1.12 | 13 | 35 | 364 | 54.6 | 596 | 332 | 1.13 | 6 |
| 12 | 22 | 380 | 48.9 | 658 | 347 | 1.13 | 0 | 13 | 380 | 70.4 | 769 | 579 | 1.20 | 23 | 35 | 380 | 56.8 | 699 | 442 | 1.16 | 9 |
| 13 | 20 | 393 | 62.6 | 745 | 436 | 1.10 | 0 | 12 | 384 | 47.6 | 665 | 246 | 1.13 | 8 | 32 | 389 | 56.8 | 715 | 374 | 1.11 | 3 |
| 14 | 5 | 403 | 49.8 | 810 | 317 | 1.18 | 0 | 12 | 435 | 31.5 | 935 | 257 | 1.12 | 33 | 17 | 426 | 39.0 | 899 | 272 | 1.14 | 24 |
| 15 | 2 | 522 | 37.5 | 1563 | 336 | 1.09 | 50 | 6 | 435 | 55.1 | 929 | 471 | 1.04 | 33 | 8 | 457 | 63.0 | 1088 | 510 | 1.06 | 38 |
| 16 | 3 | 486 | 62.6 | 1233 | 461 | 1.04 | 0 | 2 | 473 | 17.0 | 1288 | 265 | 1.21 | 50 | 5 | 481 | 45.6 | 1255 | 353 | 1.11 | 20 |
| 17 | 2 | 439 | 4.9 | 900 | 0 | 1.07 | 0 | - | - | - | - | - | - | - | 2 | 439 | 4.9 | 900 | 0 | 1.07 | 0 |
| 18 | 1 | 520 | - | 1800 | - | 1.28 | 100 | - | - | - | - | - | - | - | 1 | 520 | - | 1800 | - | 1.28 | 100 |
| 20 | - | - | - | - | - | - | - | 1 | 485 | - | 1300 | - | 1.14 | 100 | 1 | 485 | - | 1300 | - | 1.14 | 100 |
| TOTAL | 119 | | | | | | | 104 | | | | | | | 224 | | | | | | |
| MEAN | | 378 | 66 | 682 | 391 | 1.13 | | | 365 | 72.9 | 625 | 392 | 1.12 | | | 371 | 70.1 | 653 | 392 | 1.13 | |
| MEAN AGE | 11.3 | | | | | | | | | | | | | | | | | | | | |

Appendix 9. Summary of biological data by length interval for Arctic charr taken by subsistence gillnets (89 mm mesh size) in the Sylvia Grinnell River, 1977.

| LENGTH INTERVAL (MM) | MALES | | | | | | FEMALES | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | | % |
| | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT |
| 200 | 1 | 246 | 175 | - | 1.18 | 0 | 1 | 200 | 100 | - | 1.25 | 0 | 2 | 223 | 138 | 53 | 1.21 | 0 |
| 250 | 2 | 271 | 213 | 18 | 1.07 | 0 | 3 | 278 | 250 | 43 | 1.16 | 0 | 6 | 279 | 238 | 34 | 1.09 | 0 |
| 300 | 4 | 311 | 338 | 60 | 1.11 | 0 | 1 | 332 | 400 | - | 1.09 | 0 | 7 | 316 | 336 | 56 | 1.06 | 0 |
| 350 | 9 | 383 | 650 | 59 | 1.15 | 0 | 5 | 377 | 625 | 53 | 1.18 | 0 | 14 | 381 | 641 | 56 | 1.16 | 0 |
| 400 | 5 | 415 | 805 | 122 | 1.12 | 0 | 3 | 440 | 958 | 80 | 1.13 | 100 | 12 | 426 | 860 | 115 | 1.11 | 25 |
| 450 | 4 | 476 | 1125 | 65 | 1.05 | 0 | 1 | 458 | 1000 | - | 1.04 | 100 | 8 | 468 | 1094 | 91 | 1.07 | 13 |
| 500 | 1 | 530 | 1550 | - | 1.04 | 100 | 1 | 542 | 1850 | - | 1.16 | 100 | 4 | 527 | 1700 | 173 | 1.16 | 50 |
| TOTAL | 26 | | | | | | 15 | | | | | | 53 | | | | | |
| MEAN | | 384 | 688 | 341 | 1.11 | | | 371 | 673 | 440 | 1.15 | | | 389 | 734 | 410 | 1.12 | |

Appendix 10. Summary of biological data by age for Arctic charr taken by subsistence gillnets in the Sylvia Grinnell River, 1977.

| AGE (YR) | MALES | | | | | | | FEMALES | | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % |
| | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT |
| 7 | 3 | 276 | 31.6 | 233 | 80 | 1.09 | 0 | - | - | - | - | - | - | - | 3 | 276 | 31.6 | 233 | 80 | 1.09 | 0 |
| 8 | 1 | 480 | - | 1100 | - | 0.99 | 0 | 2 | 270 | 7.1 | 225 | 0 | 1.15 | 0 | 3 | 340 | 121.3 | 517 | 505 | 1.10 | 0 |
| 9 | 2 | 286 | 23.3 | 263 | 53 | 1.12 | 0 | 1 | 359 | - | 625 | - | 1.35 | 0 | 3 | 310 | 45.5 | 383 | 213 | 1.20 | 0 |
| 10 | 4 | 378 | 37.6 | 619 | 133 | 1.14 | 0 | 3 | 374 | 72.4 | 642 | 317 | 1.16 | 33 | 7 | 376 | 49.6 | 629 | 206 | 1.15 | 14 |
| 11 | 6 | 405 | 73.7 | 800 | 431 | 1.11 | 17 | 1 | 354 | - | 575 | - | 1.30 | 0 | 7 | 398 | 70.1 | 768 | 403 | 1.14 | 14 |
| 12 | 3 | 433 | 47.4 | 917 | 208 | 1.13 | 0 | 2 | 449 | 13.4 | 1025 | 35 | 1.14 | 100 | 5 | 439 | 35.1 | 960 | 160 | 1.13 | 40 |
| 13 | - | - | - | - | - | - | - | 2 | 388 | 2.8 | 613 | 53 | 1.05 | 0 | 2 | 388 | 2.8 | 613 | 53 | 1.05 | 0 |
| 14 | 1 | 385 | - | 650 | - | 1.14 | 0 | - | - | - | - | - | - | - | 1 | 385 | - | 650 | - | 1.14 | 0 |
| 16 | - | - | - | - | - | - | - | 1 | 542 | - | 1850 | - | 1.16 | 100 | 1 | 542 | - | 1850 | - | 1.16 | 100 |
| 18 | 1 | 404 | - | 725 | - | 1.10 | 0 | - | - | - | - | - | - | - | 1 | 404 | - | 725 | - | 1.10 | 0 |
| 20 | - | - | - | - | - | - | - | 1 | 448 | - | 900 | - | 1.00 | 100 | 1 | 448 | - | 900 | - | 1.00 | 100 |
| TOTAL | 21 | | | | | | | 13 | | | | | | | 34 | | | | | | |
| MEAN | | 377 | 75 | 654 | 351 | 1.11 | | | 387 | 79.5 | 738 | 434 | 1.15 | | | 381 | 75.5 | 686 | 381 | 1.13 | |
| MEAN AGE | 11.0 | | | | | | | | | | | | | | | | | | | | |

Appendix 11. Summary of biological data by length interval for Arctic charr taken by snagging in the Sylvia Grinnell River, 1977.

| LENGTH INTERVAL (MM) | MALES | | | | | | FEMALES | | | | | | COMBINED | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | | % | | LENGTH(MM) | WEIGHT(G) | | % |
| | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT | N | MEAN | MEAN | SD | K | MAT |
| 150 | - | - | - | - | - | - | 1 | 195 | 50 | - | 0.67 | 0 | 5 | 189 | 65 | 14 | 0.97 | 0 |
| 200 | 1 | 220 | 125 | - | 1.17 | 0 | 1 | 237 | 125 | - | 0.94 | 0 | 16 | 228 | 120 | 28 | 1.00 | 0 |
| 250 | 4 | 290 | 231 | 13 | 0.95 | 0 | 7 | 284 | 236 | 50 | 1.02 | 0 | 40 | 278 | 218 | 37 | 1.01 | 0 |
| 300 | 31 | 327 | 348 | 62 | 0.98 | 0 | 13 | 325 | 356 | 59 | 1.03 | 0 | 146 | 326 | 355 | 60 | 1.02 | 0 |
| 350 | 8 | 367 | 531 | 72 | 1.07 | 0 | 11 | 371 | 518 | 85 | 1.01 | 0 | 69 | 371 | 541 | 70 | 1.06 | 0 |
| 400 | 2 | 414 | 738 | 124 | 1.04 | 0 | 1 | 412 | 725 | - | 1.04 | 0 | 12 | 418 | 758 | 91 | 1.04 | 0 |
| 450 | - | - | - | - | - | - | - | - | - | - | - | - | 3 | 472 | 1158 | 153 | 1.10 | 0 |
| 500 | - | - | - | - | - | - | - | - | - | - | - | - | 1 | 508 | 1500 | - | 1.14 | 0 |
| TOTAL | 46 | | | | | | 34 | | | | | | 292 | | | | | |
| MEAN | | 332 | 382 | 133 | 1.00 | | | 328 | 379 | 154 | 1.01 | | | 328 | 391 | 188 | 1.03 | |

Appendix 12. Summary of biological data by age for Arctic charr taken by snagging in the Sylvia Grinnell River, 1977.

| AGE (YR) | MALES | | | | | | | FEMALES | | | | | | | COMBINED | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % | | LENGTH(MM) | | WEIGHT(G) | | | % |
| | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT | N | MEAN | SD | MEAN | SD | K | MAT |
| 6 | - | - | - | - | - | - | - | 2 | 246 | 72.1 | 188 | 194 | 0.96 | 0 | 2 | 246 | 72.1 | 188 | 194 | 0.96 | 0 |
| 7 | 1 | 220 | - | 125 | - | 1.17 | 0 | 4 | 307 | 32.1 | 306 | 88 | 1.04 | 0 | 5 | 289 | 47.7 | 270 | 111 | 1.07 | 0 |
| 8 | 4 | 322 | 24.1 | 325 | 89 | 0.95 | 0 | - | - | - | - | - | - | - | 4 | 322 | 24.1 | 325 | 89 | 0.95 | 0 |
| 9 | 10 | 342 | 36.4 | 425 | 152 | 1.03 | 0 | 8 | 305 | 28.1 | 284 | 82 | 0.98 | 0 | 18 | 325 | 37.2 | 363 | 142 | 1.01 | 0 |
| 10 | 6 | 323 | 19.0 | 342 | 97 | 0.99 | 0 | 7 | 354 | 19.4 | 479 | 97 | 1.07 | 0 | 13 | 340 | 24.2 | 415 | 117 | 1.03 | 0 |
| 11 | 8 | 322 | 21.0 | 331 | 66 | 0.98 | 0 | 3 | 375 | 47.6 | 567 | 213 | 1.04 | 0 | 11 | 336 | 37.0 | 395 | 156 | 1.00 | 0 |
| 12 | 7 | 349 | 26.1 | 432 | 104 | 1.00 | 0 | 5 | 352 | 21.9 | 435 | 74 | 0.99 | 0 | 12 | 350 | 23.5 | 433 | 89 | 1.00 | 0 |
| 13 | - | - | - | - | - | - | - | 1 | 375 | - | 525 | - | 1.00 | 0 | 1 | 375 | - | 525 | - | 1.00 | 0 |
| 14 | 1 | 363 | - | 550 | - | 1.15 | 0 | - | - | - | - | - | - | - | 1 | 363 | - | 550 | - | 1.15 | 0 |
| 15 | 2 | 386 | 50.9 | 638 | 265 | 1.07 | 0 | - | - | - | - | - | - | - | 2 | 386 | 50.9 | 638 | 265 | 1.07 | 0 |
| 16 | - | - | - | - | - | - | - | 1 | 398 | - | 575 | - | 0.91 | 0 | 1 | 398 | - | 575 | - | 0.91 | 0 |
| TOTAL | 39 | | | | | | | 31 | | | | | | | 70 | | | | | | |
| MEAN | | 334 | 36 | 390 | 140 | 1.01 | | | 332 | 45.8 | 394 | 151 | 1.01 | | | 333 | 40.3 | 392 | 144 | 1.01 | |
| MEAN AGE | 10.1 | | | | | | | | | | | | | | | | | | | | |

Appendix 13. Description of the relative stages of maturity of Arctic charr based on gross examination of the gonads.

| Maturity (Male and Female) | Description of Gonads |
|---|---|
| Immature | Virgin fish; gonads thin and threadlike, often incomplete. |
| Maturing | Virgin or non-virgin fish not spawning in the current year; gonads full length, firm; eggs small; gonads partially fill body cavity. |
| Mature | Fish spawning in current year; gonads full size, filling body cavity; eggs prominent, full size. |
| Ripe | Mature fish in spawning condition; eggs translucent; milt or eggs expelled under slight pressure. |
| Spent | Mature fish, completed spawning; gonads collapsed with ruptured blood vessels prominent; residual eggs present. |

Printed in Canada by
Supply and Services Canada
Printing Office
for exclusive distribution by
Fisheries and Oceans
Freshwater Institute
Winnipeg, Manitoba
Canada